

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

Journal of Algebra 284 (2005) 857–868

JOURNAL OF
Algebrawww.elsevier.com/locate/jalgebraOn the quiver of tilting modules[☆]Dieter Happel^{a,*}, Luise Unger^b^a *Fakultät für Mathematik, Technische Universität Chemnitz, D-09107 Chemnitz, Germany*^b *Fachbereich für Mathematik, Fernuniversität Hagen, D-58084 Hagen, Germany*

Received 21 September 2004

Communicated by Kent R. Fuller

Dedicated to Claus Michael Ringel on the occasion of his 60th birthday

Let Λ be an Artin algebra over a commutative Artin ring R and let $\text{mod } \Lambda$ be the category of finitely generated left Λ -modules. For a module $M \in \text{mod } \Lambda$, we denote by $\text{pd}_\Lambda M$ the projective dimension of M and by $\text{gl.dim } \Lambda$ the global dimension of Λ . A module $T \in \text{mod } \Lambda$ is called a tilting module provided the following three conditions are satisfied:

- (i) $\text{pd}_\Lambda T < \infty$,
- (ii) $\text{Ext}_\Lambda^i(T, T) = 0$ for all $i > 0$, and
- (iii) there exists an exact sequence $0 \rightarrow {}_\Lambda \Lambda \rightarrow T^0 \rightarrow \dots \rightarrow T^r \rightarrow 0$ with $T^i \in \text{add } T$ for all $0 \leq i \leq r$, where $\text{add } T$ is the full subcategory of $\text{mod } \Lambda$ whose objects are direct sums of direct summands of T .

We will say that a tilting module is basic if in a direct sum decomposition of T the indecomposable summands of T occur with multiplicity one. Unless stated otherwise all tilting modules considered here will be assumed to be basic.

[☆] The main results presented here were obtained while the two authors were visiting Beijing Normal University. The first author was supported by the EU–Asia link programme. Both authors thank their host Changchang Xi for the hospitality.

* Corresponding author.

E-mail addresses: happel@mathematik.tu-chemnitz.de (D. Happel), luise.unger@fernuni-hagen.de (L. Unger).

Following [AR] we consider for a tilting module the right perpendicular category

$$T^\perp = \{X \in \text{mod } \Lambda \mid \text{Ext}_\Lambda^i(T, X) = 0 \text{ for all } i > 0\}.$$

Important properties of these categories will be recalled in Section 1. We consider the set \mathcal{T}_Λ of all tilting modules over Λ up to isomorphism. In [HU4] the following partial order \leq on \mathcal{T}_Λ was investigated. For $T, T' \in \mathcal{T}_\Lambda$, we set $T \leq T'$ provided $T^\perp \subset T'^\perp$. One of the main results in [HU4] was that the Hasse quiver $\tilde{\mathcal{K}}_\Lambda$ of this poset allows the following description. The vertices of $\tilde{\mathcal{K}}_\Lambda$ are the elements of \mathcal{T}_Λ . There is an arrow $T' \rightarrow T$ if $T' = M \oplus X$, $T = \tilde{M} \oplus Y$ with X, Y indecomposable, and there exists a short exact sequence $0 \rightarrow X \rightarrow \tilde{M} \rightarrow Y \rightarrow 0$ with $\tilde{M} \in \text{add } M$. In Section 2 we will recall the reason behind this. Moreover, we will show the following for a representation-finite algebra Λ of finite global dimension d : For each $0 \leq i \leq d$ there exists a tilting module $T \in \tilde{\mathcal{K}}_\Lambda$ with $\text{pd}_\Lambda T = i$.

The main aim of this article is to investigate the local structure of $\tilde{\mathcal{K}}_\Lambda$ in more detail. In particular, we will establish a precise relationship between the number of neighbours of a vertex $T \in \tilde{\mathcal{K}}_\Lambda$ and the add T -coresolution of ${}_\Lambda \Lambda$ in the definition of a tilting module. We will also discuss the number and length of paths leading or starting in a given vertex. It is interesting to point out that the finiteness of certain paths in $\tilde{\mathcal{K}}_\Lambda$ starting in a vertex T is related to the generalised Nakayama conjecture. For details we refer to Section 4. In the special case of hereditary algebras we can say more. In this case we say that a vertex $T \in \tilde{\mathcal{K}}_\Lambda$ is saturated if the number of neighbours coincides with the number n of distinct simple modules. It is easy to see that the number of neighbours is always bounded by n . We will show that a vertex T is saturated if and only if each coordinate of the dimension vector of T is at least two. The results for hereditary algebras are contained in Section 3.

We denote the composition of morphisms $f: X \rightarrow Y$ and $g: Y \rightarrow Z$ in a given category \mathcal{K} by fg .

The notation and terminology introduced here will be fixed throughout this article. For unexplained representation-theoretic terminology, we refer to [R] or [ARS].

1. Preliminaries

In this section we will briefly recall the basic definitions and results from tilting theory we will use in the main part of this article.

Given any Λ -module Z , we may decompose it: $Z \simeq \bigoplus_{i=1}^m Z_i^{d_i}$, where Z_i is indecomposable and $d_i > 0$ for all i and moreover $Z_i \not\simeq Z_j$ for $i \neq j$. We call Z basic in case $d_i = 1$ for all i . The uniquely determined number m occurring in the direct sum decomposition above will be denoted by $\delta(Z)$. It is not hard to see that a tilting module T satisfies $\delta(T) = n = \text{rk } K_0(\Lambda)$, where $K_0(\Lambda)$ denotes the Grothendieck group of Λ . We say that a module M is a partial tilting module if M is a direct summand of a tilting module. A partial tilting module M with $\delta(M) = n - 1$ is called an almost complete partial tilting module. Given a partial tilting module M , we say that C is a complement to M if $M \oplus C$ is a tilting module and $\text{add } M \cap \text{add } C = 0$. Note that a complement to an almost complete partial tilting module is indecomposable. We will see later that special complements are important.

Let M be a partial tilting module and C be a complement to M , then C is called a source complement if

$$(M \oplus C)^\perp = M^\perp.$$

We refer to [HU1, HU3] for a characterisation and the concept of sink complements.

More generally, we call $X \in \text{mod } \Lambda$ self-orthogonal if $\text{Ext}_\Lambda^i(X, X) = 0$ for all $i > 0$, and we call X exceptional if X is self-orthogonal and satisfies $\text{pd}_\Lambda X < \infty$. For more details, we refer to [H]. So partial tilting modules are typical examples of exceptional modules.

Let $T = \bigoplus_{j=1}^n T_j$ be a tilting module over Λ . For $1 \leq i \leq n$, we define $T[i] = \bigoplus_{j \neq i} T_j$. Note that $T[i]$ is an almost complete partial tilting module.

For $X \in \text{mod } \Lambda$, we denote by $\text{fac } X$ the full subcategory of $\text{mod } \Lambda$ consisting of those modules Y which are epimorphic images of modules in $\text{add } X$. Dually, we denote by $\text{sub } X$ the full subcategory of $\text{mod } \Lambda$ consisting of submodules of modules in $\text{add } X$.

Let ${}_A T$ be a tilting module and let $X \in T^\perp$. As shown, for example, in [CHU], we have that $X \in \text{fac } T$. So there is a short exact sequence $0 \rightarrow K_0 \rightarrow T_0 \rightarrow X \rightarrow 0$ with $T_0 \in \text{add } T$ and $K_0 \in T^\perp$. Inductively, we obtain a long exact sequence

$$\cdots \rightarrow T_r \xrightarrow{f_r} T_{r-1} \rightarrow \cdots \rightarrow T_0 \xrightarrow{f_0} X \rightarrow 0 \quad (*)$$

with $T_i \in \text{add } T$ and $K_i = \ker f_i \in T^\perp$ for all $i \geq 0$. We call $(*)$ an $\text{add } T$ -resolution of X , and if $(*)$ is minimal, a minimal $\text{add } T$ -resolution. If $\text{pd}_\Lambda X < \infty$, then X admits a finite $\text{add } T$ -resolution $0 \rightarrow T_r \rightarrow T_{r-1} \rightarrow \cdots \rightarrow T_0 \rightarrow X \rightarrow 0$. The following lemma is shown in [HU4] and will be needed in Section 3.

Lemma 1.1. *Let ${}_A T$ be a tilting module. Let $X \in T^\perp$ be an exceptional module with $X \notin \text{add } T$. If $0 \rightarrow T_r \rightarrow T_{r-1} \rightarrow \cdots \rightarrow T_0 \rightarrow X \rightarrow 0$ is a minimal $\text{add } T$ -resolution of X , then $\text{add } T_r \cap \text{add } T_0 = 0$.*

We recall from [HU3] the structure of complements to almost complete partial tilting modules. This will be needed in section three. We first recall the following proposition.

Proposition 1.2. *Let M be an almost complete tilting module. Let Y be an indecomposable complement generated by M . Then*

- (1) M is faithful;
- (2) there exists an indecomposable complement X not isomorphic to Y ;
- (3) there exists an exact sequence $0 \rightarrow X \rightarrow E \rightarrow Y \rightarrow 0$ with $E \in \text{add } M$;
- (4) $\text{Ext}_\Lambda^i(X, Y) = 0$ for $i > 0$ and $\text{Ext}_\Lambda^i(Y, X) = 0$ for $i > 1$; and
- (5) X is uniquely determined by the property (3).

We also point out that Y is uniquely determined by the property (3). Note that in (3) E is the minimal right $\text{add } M$ -approximation of Y , as well as the minimal left $\text{add } M$ -approximation of X . The exact sequence in (3) is called the connecting sequence.

The following is easily deduced.

Corollary 1.3. *Let M be an almost complete partial tilting module. Then M admits a source complement X_0 . More precisely, for each complement Y to M with $Y \in \text{fac } M$ there exists a long exact sequence*

$$0 \rightarrow X_0 \rightarrow M^1 \xrightarrow{f_1} M^2 \rightarrow \cdots \rightarrow M^{r-1} \xrightarrow{f_{r-1}} M^r \xrightarrow{f_r} X_r = Y \rightarrow 0$$

with $\ker f_i = X_{i-1}$ for $1 \leq i \leq r$, $M^i \in \text{add } M$ for $1 \leq i \leq r$ and X_0, \dots, X_r complements to M where $r \leq \text{pd}_A Y$.

In the following, we will always denote by X_0 the source complement to an almost complete partial tilting module M . The finitistic dimension of Λ is defined as

$$\text{fd}(\Lambda) = \sup\{\text{pd}_\Lambda X \mid X \in \mathcal{P}^{<\infty}(\Lambda)\}.$$

Theorem 1.4. *Let Λ be an Artin algebra. Let M be an almost complete partial tilting module. Then M has at most finitely many non-isomorphic complements if and only if M admits a sink complement. If there are $r + 1$ non-isomorphic complements for some $r \geq 1$, then $r \leq \text{fd}(\Lambda)$ and there exists a long exact sequence*

$$0 \rightarrow X_0 \rightarrow M^1 \xrightarrow{f_1} M^2 \rightarrow \cdots \rightarrow M^{r-1} \xrightarrow{f_{r-1}} M^r \xrightarrow{f_r} X_r \rightarrow 0$$

with $\ker f_i = X_{i-1}$ for $1 \leq i \leq r$, $M^i \in \text{add } M$ for $1 \leq i \leq r$ and X_0, \dots, X_r complements to M . In particular, if $\text{fd}(\Lambda) < \infty$, then M has finitely many complements.

Let M be an almost complete partial tilting module and X a complement to M . Then $X \simeq X_s$ for some s . Then we will say that the number of complements to M preceding X is $s + 1$. If there is a finite number $r + 1$ of complements to M , we say that the number of complements to M succeeding X is $r + 1 - s$. Otherwise, we will say that there are infinitely many complements to M succeeding X .

We will also point out the main connection to the generalised Nakayama conjecture but refer for the details to [HU3].

Theorem 1.5. *Let M be an almost complete partial tilting module with source complement X_0 . Let $\Gamma = \text{End}_\Lambda(M \oplus X_0)$. Let ${}_\Gamma S$ be the simple top of the indecomposable projective Γ -module $\text{Hom}_\Lambda(M \oplus X_0, X_0)$. If M admits t complements different from X_0 , then $\text{Ext}_\Gamma^j(D(\Gamma_\Gamma), S) = 0$ for $0 \leq j < t$.*

2. Hasse quiver and consequences

In the introduction we have defined the partially ordered set $(\mathcal{T}_\Lambda, \leq)$ of tilting modules of an Artin algebra Λ . We start by observing some elementary properties. The assertion 2.1(a) is contained in [HU4].

Lemma 2.1. *Let Λ be an Artin algebra.*

- (a) $T \leq T'$ in \mathcal{T}_Λ if and only if $T \in T'^\perp$.
- (b) If $T \leq T'$ in \mathcal{T}_Λ then $\text{pd}_\Lambda T \geq \text{pd}_\Lambda T'$.

Proof. If $T \leq T'$ then $T \in T'^\perp$. Let $r = \text{pd}_\Lambda T$. Then by the defining property of a tilting module there is an exact sequence

$$0 \rightarrow {}_\Lambda \Lambda \rightarrow T^0 \rightarrow \cdots \rightarrow T^s \rightarrow 0$$

with $T^i \in \text{add } T$ for all $0 \leq i \leq s$. It is well known that we may choose such a sequence with $s = r$. Since $T \in T'^\perp$, it follows that $\text{Ext}_\Lambda^{i+1}(T', {}_\Lambda \Lambda) = 0$ for $i \geq r$. Since $\text{pd}_\Lambda T' < \infty$, we infer that $\text{pd}_\Lambda T' \leq r$. \square

The following theorem and its corollary are contained in [HU4] and they are stated for the convenience of the reader. The theorem will be used in the next section.

Theorem 2.2. *Let $T, T' \in \mathcal{T}_\Lambda$ with $T \leq T'$ and $T \neq T'$. Then there exists $T'' \in \mathcal{T}_\Lambda$ with $T' \rightarrow T''$ in $\vec{\mathcal{K}}_\Lambda$ such that $T \leq T''$. In particular, $\vec{\mathcal{K}}_\Lambda$ is the Hasse quiver of \mathcal{T}_Λ .*

Corollary 2.3. *If $\vec{\mathcal{K}}_\Lambda$ has a finite connected component \mathcal{C} , then $\vec{\mathcal{K}}_\Lambda = \mathcal{C}$.*

We will deduce some further consequences.

Corollary 2.4. *If $\vec{\mathcal{K}}_\Lambda$ has a finite component \mathcal{C} , let T be the minimal element in \mathcal{C} . Let $d = \text{pd}_\Lambda T$. Then for each $0 \leq i \leq d$ there exists a tilting module T_i with $\text{pd}_\Lambda T_i = i$.*

Proof. By Corollary 2.3 we have that $\mathcal{C} = \vec{\mathcal{K}}_\Lambda$. If $T' \rightarrow T''$ is an arrow in $\vec{\mathcal{K}}_\Lambda$, we have by Lemma 2.1 that $\text{pd}_\Lambda T'' \geq \text{pd}_\Lambda T'$. By the definition of arrows in $\vec{\mathcal{K}}_\Lambda$, we actually have that

$${}_ \Lambda T' \leq \text{pd}_\Lambda T'' \leq \text{pd}_\Lambda T' + 1. \quad (*)$$

Since $\vec{\mathcal{K}}_\Lambda$ is finite there is a path in $\vec{\mathcal{K}}_\Lambda$ from the unique source ${}_\Lambda \Lambda$ of $\vec{\mathcal{K}}_\Lambda$ to the unique minimal element T of $\vec{\mathcal{K}}_\Lambda$ of the form

$${}_\Lambda \Lambda \rightarrow T_0 \rightarrow T_1 \rightarrow \cdots \rightarrow T_{s-1} \rightarrow T_s \rightarrow T.$$

Now $\text{pd}_\Lambda {}_\Lambda \Lambda = 0$ and $\text{pd}_\Lambda T = d$. Using the chain (*) of inequalities shows the assertion. \square

The following is an immediate consequence of the previous corollary.

Corollary 2.5. *Let Λ be a representation-finite Artin algebra of finitistic dimension d . Then for each $0 \leq i \leq d$ there exists an indecomposable exceptional module E_i with $\text{pd}_\Lambda E_i = i$.*

Proof. Let T be the sink in $\vec{\mathcal{K}}_\Lambda$. It follows from [HU2,HU4] that each Λ -module X with $\text{pd } X < \infty$ satisfies $X \in \text{sub } T$. Since the finitistic dimension of Λ is d there is X with $\text{pd } X = d$. So let

$$0 \rightarrow X \rightarrow T' \rightarrow Y \rightarrow 0$$

be exact with $T' \in \text{add } T$. Since $\text{pd } X = d$ and $\text{pd } Y \leq d$, we infer that $\text{pd } T' = d$. So the assertion follows from Corollary 2.4. \square

We refer to [HU4] for a result about the existence of minimal elements in the partial order \mathcal{T}_Λ which uses [HU2].

3. Local structure of the quiver for hereditary algebras

In the introduction we have defined the partially ordered set $(\mathcal{T}_\Lambda, \leq)$ and the quiver $\vec{\mathcal{K}}_\Lambda$ of tilting modules. In this section we will mainly be interested in the local structure of $\vec{\mathcal{K}}_\Lambda$ for hereditary algebras. So for the rest of this section, Λ denotes a hereditary Artin algebra.

Let $T \in \vec{\mathcal{K}}_\Lambda$. We denote by $s(T)$ (respectively $e(T)$) the number of arrows starting (respectively ending) at T in $\vec{\mathcal{K}}_\Lambda$ (see Fig. 1).

Lemma 3.1. $s(T) + e(T) \leq \text{rk } K_0(\Lambda) = n$.

Proof. Let $T = \bigoplus_{i=1}^n T_i \in \vec{\mathcal{K}}_\Lambda$. For each $1 \leq i \leq n$, we consider the almost complete partial tilting module $T[i]$. For $T[i]$, we have an indecomposable complement T_i . If T_i is generated by $T[i]$, we obtain from Proposition 1.2 a complement X_i which is cogenerated by $T[i]$, hence an arrow from $T[i] \oplus X_i \rightarrow T$ in $\vec{\mathcal{K}}_\Lambda$. If T_i is cogenerated by $T[i]$, we obtain from Proposition 1.2 a complement Y_i which is generated by $T[i]$, hence an arrow from $T \rightarrow T[i] \oplus Y_i$ in $\vec{\mathcal{K}}_\Lambda$. Thus to show the assertion, it is enough to show that T_i cannot be both generated and cogenerated by $T[i]$. If T_i is both generated and cogenerated by $T[i]$, we obtain that $T[i]$ admits at least three non-isomorphic complements. Since Λ is hereditary, this contradicts Theorem 1.4. \square

We say that T is saturated if equality holds. Note that the proof of Lemma 3.1 shows that T is saturated if and only if $T[i]$ is faithful or equivalently sincere for each $1 \leq i \leq n$.




Fig. 1.

For a Λ -module X we denote by $\mathbf{dim} X \in K_0(\Lambda)$ its dimension vector which we will identify with the corresponding element in \mathbb{Z}^n .

Proposition 3.2. $T \in \vec{\mathcal{K}}_\Lambda$ is saturated iff $(\mathbf{dim} T)_i \geq 2$ for all $1 \leq i \leq n$.

Proof. If there is $1 \leq i \leq n$ with $(\mathbf{dim} T)_i = 1$, choose an indecomposable direct summand X of T with $(\mathbf{dim} X)_i = 1$. Let $T = M \oplus X$. Then $(\mathbf{dim} M)_i = 0$, so M is not sincere. Thus T is not saturated by the remark following the proof of Lemma 3.1.

Conversely, assume that $T \in \vec{\mathcal{K}}_\Lambda$ is not saturated. So there is an indecomposable direct summand X of $T = M \oplus X$ such that M is not sincere. So there exists an indecomposable injective Λ -module I such that $\mathrm{Hom}_\Lambda(M, I) = 0$. Let $\Gamma = \mathrm{End}_\Lambda T$. Then $\mathrm{Hom}_\Lambda(T, I)$ is an indecomposable Γ -module. Set $P_X = \mathrm{Hom}_\Lambda(T, X)$ and $S_X = \mathrm{top} P_X$. Now

$$\mathrm{Hom}_\Lambda(T, I) \simeq S_X^{\dim \mathrm{Hom}_\Lambda(T, I)} \simeq S_X^{\dim \mathrm{Hom}_\Lambda(X, I)}.$$

Since $\mathrm{Hom}_\Lambda(T, I)$ is indecomposable, we infer that $\dim \mathrm{Hom}_\Lambda(T, I) = 1$. Thus there is $1 \leq i \leq n$ with $(\mathbf{dim} T)_i = 1$. This finishes the proof. \square

We have the following consequence for the path algebra Λ of $\Delta = \mathbb{A}_n$ with linear orientation over a field k . There are no saturated vertices in $\vec{\mathcal{K}}_\Lambda$. In fact, it is well known that for a tilting module T over the path algebra of \mathbb{A}_n with linear orientation the unique indecomposable projective–injective module is a direct summand and the direct sum of the other indecomposable summands is not sincere. Thus there is a coordinate of the dimension vector of T with value 1.

We will point out a different way of detecting arrows in $\vec{\mathcal{K}}_\Lambda$. For this, let $T \in \mathcal{T}_\Lambda$ and $\Gamma = \mathrm{End} T$. Let $(\mathcal{X}(T), \mathcal{Y}(T))$ be the torsion pair in $\mathrm{mod} \Gamma$ induced by T (for details, compare [HR]). Let $T = M \oplus X$ and let P_X, S_X be as above.

Proposition 3.3. *With the above notation the following hold.*

- (1) $S_X \in \mathcal{X}(T)$ if and only if $X \in \mathrm{fac} M$.
- (2) $X \in \mathrm{sub} M$ if and only if $S_X \in \mathcal{Y}(T)$ and $T \otimes_\Gamma S_X$ is not injective.

Proof. (1) If $X \in \mathrm{fac} M$, we consider an exact sequence $0 \rightarrow Y \rightarrow \tilde{M} \rightarrow X \rightarrow 0$ with $\tilde{M} \rightarrow X$ a minimal right add M -approximation of X . Applying $\mathrm{Hom}_\Lambda(T, -)$ to this sequence yields an exact sequence of Γ -modules

$$\mathrm{Hom}_\Lambda(T, \tilde{M}) \rightarrow P_X \rightarrow \mathrm{Ext}_\Lambda^1(T, Y) \rightarrow 0.$$

Now $\mathrm{Ext}_\Lambda^1(T, Y) \simeq S_X$ and $\mathrm{Ext}_\Lambda^1(T, Y) \in \mathcal{X}(T)$ by tilting theory, thus $S_X \in \mathcal{X}(T)$.

Conversely, assume that $S_X \in \mathcal{X}(T)$. Consider the exact sequence of Γ -modules

$$0 \rightarrow K_X \rightarrow P_X \rightarrow S_X \rightarrow 0.$$

Since $P_X \in \mathcal{Y}(T)$, we see that $K_X \in \mathcal{Y}(T)$, for $\mathcal{Y}(T)$ is a torsion-free class. Now $\mathrm{End} X$ is a division ring, since X is an indecomposable direct summand of a tilting module. By

tilting theory, we infer that the same holds for $\text{End } P_X$. Hence S_X is not a composition factor of K_X and so we see that $K_X \in \text{fac Hom}_\Lambda(T, M)$. But this implies that $X \in \text{fac } M$.

(2) If $X \in \text{sub } M$, we consider an exact sequence

$$0 \rightarrow X \rightarrow \tilde{M} \rightarrow Y \rightarrow 0$$

with $X \rightarrow \tilde{M}$ a minimal left add M -approximation of X . Applying $\text{Hom}_\Lambda(T, -)$ to this sequence yields an exact sequence of Γ -modules

$$0 \rightarrow P_X \rightarrow \text{Hom}_\Lambda(T, \tilde{M}) \rightarrow \text{Hom}_\Lambda(T, Y) \rightarrow 0.$$

Applying $\text{Hom}_\Gamma(-, S_X)$ to this exact sequence shows that $\text{Ext}_\Gamma^1(\text{Hom}_\Lambda(T, Y), S_X) \neq 0$. Since $\text{Hom}_\Lambda(T, Y) \in \mathcal{Y}(T)$ and $(\mathcal{X}(T), \mathcal{Y}(T))$ is a split torsion pair, we infer that $S_X \in \mathcal{Y}(T)$. Since $X \in \text{sub } M$, we have that M is sincere. Assume to the contrary that $T \otimes_\Gamma S_X = I$ is an injective Λ -module. Then $\text{Hom}_\Lambda(T, I) = S_X$. Now $\text{Hom}_\Lambda(X, I) = \text{Hom}_\Lambda(P_X, S_X) \neq 0$ shows that $\text{Hom}_\Lambda(M, I) = 0$. Thus M is not sincere, a contradiction.

Conversely, let $S_X \in \mathcal{Y}(T)$ and $T \otimes_\Gamma S_X$ not injective as Λ -module. Assume to the contrary that $X \notin \text{sub } M$. Since $S_X \in \mathcal{Y}(T)$, we have by (1) that $X \notin \text{fac } M$. Hence M is not sincere. So there exists an indecomposable injective Λ -module I with $\text{Hom}_\Lambda(M, I) = 0$ and $\text{Hom}_\Lambda(T, I) = S_X$. But then by tilting theory we have that $T \otimes_\Gamma S_X = I$, a contradiction. \square

This yields a module theoretic characterisation of saturated vertices.

Corollary 3.4. *Let $T \in \vec{\mathcal{K}}_\Lambda$ and $\Gamma = \text{End } T$. Then T is saturated if and only if for all simple Γ -modules $S \in \mathcal{Y}(T)$ the Λ -module $T \otimes_\Gamma S$ is not injective.*

Proof. Let $T \in \vec{\mathcal{K}}_\Lambda$ and $S = S_X \in \mathcal{Y}(T)$ be a simple Γ -module. Let $T = M \oplus X$. Since $S_X \in \mathcal{Y}(T)$, we have by Proposition 3.3(1) that $X \notin \text{fac } M$. Since T is saturated, we have that $X \in \text{sub } M$. But then by Proposition 3.3(2) we have that $T \otimes_\Gamma S_X$ is not injective as Λ -module.

Conversely, let X be an indecomposable direct summand of $T = M \oplus X$. If $S_X \in \mathcal{X}(T)$, we have by Proposition 3.3(1) that $X \in \text{fac } M$. If $S_X \in \mathcal{Y}(T)$, we have by Proposition 3.3(2) that $X \in \text{sub } M$ using the assumption that $T \otimes_\Gamma S$ is not injective as Λ -module. Thus T is saturated. \square

The following theorem is contained in [U].

Theorem 3.5. *Each connected component of $\vec{\mathcal{K}}_\Lambda$ contains a non-saturated vertex.*

Remark 3.6. We conjecture that each connected component of $\vec{\mathcal{K}}_\Lambda$ contains only finitely many non-saturated vertices.

In the next section we will present a result about paths starting and ending in a given vertex $T \in \vec{\mathcal{K}}_\Lambda$. As a special case, we give here the result for neighbours. For a proof, we

refer to the next section. For this let $0 \rightarrow {}_{\Lambda}\Lambda \rightarrow T^0 \rightarrow T^1 \rightarrow 0$ and $0 \rightarrow T_1 \rightarrow T_0 \rightarrow D\Lambda_{\Lambda} \rightarrow 0$ be minimal add T -resolutions.

Proposition 3.7. $s(T) = n - \delta(T_0) = \delta(T_1)$ and $e(T) = n - \delta(T^0) = \delta(T^1)$.

Proof. The equalities $s(T) = \delta(T_1)$ and $e(T) = \delta(T^1)$ follow from the main theorem in the next section. The equalities $n - \delta(T_0) = \delta(T_1)$ and $n - \delta(T^0) = \delta(T^1)$ follow from Lemma 1.1 and its dual by observing that each indecomposable direct summand of T has to occur in the above resolutions. \square

Corollary 3.8. $T \in \vec{\mathcal{K}}_{\Lambda}$ is saturated if and only if $n = \delta(T^0) + \delta(T_0)$. In particular, ${}_{\Lambda}\Lambda$ is not saturated.

Proof. This follows immediately from Proposition 3.7. \square

We close this section by a result on regular tilting modules for wild hereditary algebras. In general, a regular tilting module will have neighbours in $\vec{\mathcal{K}}_{\Lambda}$ which have preprojective or preinjective indecomposable direct summands.

Proposition 3.9. Let Λ be a wild hereditary algebra with at least three simple modules. Let $T \in \vec{\mathcal{K}}_{\Lambda}$ be regular. Then there exist $T', T'' \in \vec{\mathcal{K}}_{\Lambda}$ regular with $T' \rightarrow T \rightarrow T''$ in $\vec{\mathcal{K}}_{\Lambda}$.

Proof. Let $T \in \vec{\mathcal{K}}_{\Lambda}$ be a regular tilting module. Then there is r_0 such that for all $r \geq r_0$ we have that $\text{Ext}_{\Lambda}^1(T, \tau^r T) = 0$ (compare, for example, [K]). Thus $\tau^r T \leq T$ in \mathcal{T}_{Λ} for all $r \geq r_0$. By Theorem 2.2 and the fact that T has only finitely many neighbours in $\vec{\mathcal{K}}_{\Lambda}$, there exists $T'' \in \vec{\mathcal{K}}_{\Lambda}$ and $T \rightarrow T''$ in $\vec{\mathcal{K}}_{\Lambda}$ such that $\tau^r T \leq T''$ for infinitely many $r \geq r_0$. Let $T = M \oplus X$ and $T'' = M \oplus Y$. The arrow in $\vec{\mathcal{K}}_{\Lambda}$ yields an exact sequence $0 \rightarrow X \rightarrow \tilde{M} \rightarrow Y \rightarrow 0$ with $\tilde{M} \in \text{add } M$. Since T is regular, we see that Y is not preprojective. Let $r \geq r_0$ with $\tau^r T \leq T''$. Then $\text{Ext}_{\Lambda}^1(Y, \tau^r T) = 0$, hence $\text{Hom}_{\Lambda}(\tau^{r-1} T, Y) = 0$. Assume that Y is preinjective, so $Y = \tau^s I$ for some $s \geq 0$ and some indecomposable injective Λ -module. So $\text{Hom}_{\Lambda}(\tau^{r-1-s} T, I) = 0$. But $\tau^t T$ is a tilting module for all t , hence sincere. This is a contradiction, so Y is regular, hence T'' is regular.

The existence of the arrow $T' \rightarrow T$ with T' regular is proved dually. \square

4. Local structure of the quiver for arbitrary algebras

In this section, let Λ be an arbitrary Artin algebra. We will investigate here certain paths starting or ending in a given vertex $T \in \vec{\mathcal{K}}_{\Lambda}$. Before proving the main theorem in this section, we will need some preparation.

Lemma 4.1. Let Λ be an Artin algebra and M an almost complete tilting module. Let X be a complement to M . Let $T = M \oplus X$. Let

$$0 \rightarrow K \rightarrow V \rightarrow D(\Lambda_{\Lambda}) \rightarrow 0 \quad (*)$$

be the minimal right add T -approximation of $D(\Lambda_A)$. Then $V \in \text{add } M$, if and only if $X \in \text{sub } M$.

Proof. Let $X \in \text{mod } \Lambda$ and $\mu : X \rightarrow D(\Lambda_A)^s$ be an injective map for some $s > 0$. Since $(*)$ is the minimal right add T -approximation of $D(\Lambda_A)$, we have that $\text{Ext}_A^1(X, K) = 0$. This implies that μ factors over $V^s \rightarrow (D(\Lambda_A))^s \rightarrow 0$, so $X \in \text{sub } V$. Thus by assumption $X \in \text{sub } M$.

Conversely, assume that $X \in \text{sub } M$. In particular, we have that M is faithful. Consider as above the connecting sequence $0 \rightarrow X \rightarrow \tilde{M} \rightarrow Y \rightarrow 0$ with $\tilde{M} \in \text{add } M$. Moreover, let

$$0 \rightarrow K' \rightarrow M' \rightarrow D(\Lambda_A) \rightarrow 0 \quad (**)$$

be the minimal right add M -approximation of $D(\Lambda_A)$. Then we have $\text{Ext}_A^i(X, K') = \text{Ext}_A^{i+1}(Y, K') = \text{Ext}_A^{i+1}(Y, M') = 0$, for $M \oplus Y$ is a tilting module. Thus $(**)$ is the minimal right add T -approximation of $D(\Lambda_A)$. So $V \in \text{add } M$. This proves the assertion.

We will also use the dual statement which we state without proof. \square

Lemma 4.2. Let Λ be an Artin algebra and M an almost complete tilting module. Let X be a complement to M . Let $T = M \oplus X$. Let

$$0 \rightarrow {}_A\Lambda \rightarrow W \rightarrow Q \rightarrow 0 \quad (*)$$

be the minimal left add T -approximation of ${}_A\Lambda$. Then $W \in \text{add } M$ if and only if $X \in \text{fac } M$.

We will now fix some more notation. For $T \in \vec{\mathcal{K}}_A$, consider the following two minimal add T -resolutions:

$$0 \rightarrow {}_A\Lambda \rightarrow T^0 \rightarrow \cdots \rightarrow T^r \rightarrow 0, \quad (+)$$

$$\cdots \rightarrow T_s \rightarrow \cdots \rightarrow T_0 \rightarrow D\Lambda_A \rightarrow 0. \quad (++)$$

Let X be an indecomposable direct summand of T . We choose $i(X)$ minimal such that X is a direct summand of $T^{i(X)}$. Note that each indecomposable direct summand X of T has to occur in $(+)$, so $i(X)$ is well defined. If X occurs in $(++)$, we choose $j(X)$ minimal such that X is a direct summand of $T_{j(X)}$. Otherwise, we set $j(X) = \infty$.

Note that the validity of the generalised Nakayama conjecture will imply that $j(X) < \infty$. For this, compare [HU3] and Theorem 1.5.

Lemma 4.3. Let T be a tilting module and let X be an indecomposable direct summand of $T = M \oplus X$. Then the number of complements to M preceding X is $i(X) + 1$.

Proof. Let

$$0 \rightarrow {}_A\Lambda \rightarrow T^0 \rightarrow \cdots \rightarrow T^r \rightarrow 0$$

be the minimal add T -resolution of ${}_A\Lambda$. We will show the assertion by induction on $i(X)$. If $i(X) = 0$, then $X \in \text{add } T^0$. By Lemma 4.2 we infer that $X \notin \text{fac } M$, hence X is the source complement to M . Since in this case there is precisely one complement to M preceding X , the assertion holds. If $i(X) > 0$, let $X = X_s$. We consider the long exact sequence of complements to M preceding X_s :

$$0 \rightarrow X_0 \rightarrow M^0 \rightarrow \cdots \rightarrow M^{s-1} \rightarrow X_s \rightarrow 0$$

with $M^i \in \text{add } M$ for $0 \leq i \leq s-1$. Moreover, let $0 \rightarrow X_{s-1} \rightarrow M^{s-1} \rightarrow X_s \rightarrow 0$ be the connecting sequence. Let $T' = M \oplus X_{s-1}$. By induction, we have that $s = i(X_{s-1}) + 1$. We consider the start of the minimal add T' -resolution of ${}_A\Lambda$:

$$0 \rightarrow {}_A\Lambda \rightarrow T'^0 \rightarrow \cdots \rightarrow T'^s. \quad (*)$$

By construction, we have that $X_{s-1} \in \text{add } T'^{s-1}$ and $T'^i \in \text{add } M$ for $0 \leq i < s-1$. From $(*)$ consider $0 \rightarrow Q^{s-2} \rightarrow T'^{s-1} \rightarrow Q^{s-1} \rightarrow 0$. Since $X_{s-1} \in \text{sub } M$, we have that $Q^{s-2} \in \text{sub } M$. Let

$$0 \rightarrow Q^{s-2} \rightarrow \tilde{M} \rightarrow Q'^{s-1} \rightarrow 0 \quad (**)$$

be the minimal left add M -approximation of Q^{s-2} . Now we have $\text{Ext}_A^i(Q'^{s-1}, X_s) = \text{Ext}_A^{i+1}(Q'^{s-1}, X_{s-1}) = \text{Ext}_A^i(Q^{s-2}, X_{s-1}) = 0$. Thus $(**)$ is also the minimal left add T -approximation of Q^{s-2} . So we have that $s \leq i(X_s)$. Assume that $s < i(X_s)$. Then $Q'^{s-1} \in \text{sub } M$ and the minimal left add M -approximation $0 \rightarrow Q'^{s-1} \rightarrow M' \rightarrow Q^s \rightarrow 0$ of Q'^{s-1} is also the minimal left add T -approximation of Q'^{s-1} . Thus $\text{Ext}_A^i(Q^s, X_s) = 0$. But $0 = \text{Ext}_A^i(Q^s, X_s) = \text{Ext}_A^{i+1}(Q^s, X_{s-1}) = \text{Ext}_A^i(Q'^{s-1}, X_{s-1})$ shows that $(**)$ is also the minimal left add T' -approximation, in contrast to $i(X_{s-1}) = s-1$. This finishes the proof of the assertion. \square

We will use the following dual lemma without proof.

Lemma 4.4. *Let T be a tilting module and let X be an indecomposable direct summand of $T = M \oplus X$. Then the number of complements to M succeeding X is $j(X) + 1$.*

Theorem 4.5. *For each indecomposable direct summand X of T , there exists a path $w(X)$ in $\vec{\mathcal{K}}_A$ of length $i(X)$ ending at T and a path $u(X)$ in $\vec{\mathcal{K}}_A$ of length $j(X)$ starting in T . These paths are pairwise disjoint.*

Proof. By Lemmas 4.3 and 4.4, the assertions on the paths follow directly from the definition of arrows in $\vec{\mathcal{K}}_A$. So it remains to show that the paths are pairwise disjoint. Let $T = M \oplus X = N \oplus Y \in \vec{\mathcal{K}}_A$ with X not isomorphic to Y . So X is an indecomposable direct summand of N and Y is an indecomposable direct summand of M . Let

$$T_{i(X)} \rightarrow \cdots \rightarrow T_1 \rightarrow M \oplus X \rightarrow T^1 \rightarrow \cdots \rightarrow T^{j(X)}$$

and

$$\overline{T}_{i(Y)} \rightarrow \cdots \rightarrow \overline{T}_1 \rightarrow N \oplus Y \rightarrow \overline{T}^1 \rightarrow \cdots \rightarrow \overline{T}^{j(Y)}$$

be the paths given by Lemmas 4.3 and 4.4. Suppose that these paths are not disjoint. Since $\vec{\mathcal{K}}_\Lambda$ does not contain oriented cycles, there exist i and j such that $T_i = \overline{T}_j$ or $T^i = \overline{T}^j$. Assume that $T_i = \overline{T}_j$. By construction, we have that $T_i = M \oplus X_i$ for some complement X_i to M and that $\overline{T}_j = N \oplus Y_j$ for some complement Y_j to N . Since X is not isomorphic to Y , the same holds for X_i and Y_j . So X_i is a direct summand of N . Thus $\text{Ext}_\Lambda^t(X, X_i) = 0$ for all $t > 0$. But by Theorem 1.4, we have that $\text{Ext}_\Lambda^t(X, X_i) \neq 0$, a contradiction. The case that $T^i = \overline{T}^j$ can be ruled out similarly. \square

References

- [AR] M. Auslander, I. Reiten, Applications of contravariantly finite subcategories, *Adv. Math.* 86 (1991) 111–152.
- [ARS] M. Auslander, I. Reiten, S. Smalø, *Representation Theory of Artin Algebras*, Cambridge University Press, 1995.
- [CHU] F. Coelho, D. Happel, L. Unger, Complements to partial tilting modules, *J. Algebra* 170 (1994) 184–205.
- [H] D. Happel, Selforthogonal modules, in: *Abelian Groups and Modules*, Kluwer Academic, pp. 257–276.
- [HU1] D. Happel, L. Unger, Partial tilting modules and covariantly finite subcategories, *Comm. Algebra* 22 (1994) 1723–1727.
- [HU2] D. Happel, L. Unger, Modules of finite projective dimensions and cocovers, *Math. Ann.* 306 (1996) 445–457.
- [HU3] D. Happel, L. Unger, Complements and the generalized Nakayama conjecture, in: *Algebras and Modules II*, Proc. ICRA VIII, Geiranger, in: CMS Conf. Proc., vol. 24, 1998, pp. 293–310.
- [HU4] D. Happel, L. Unger, On a partial order of tilting modules, *Represent. Theory*, in press.
- [HR] D. Happel, C.M. Ringel, Tilted algebras, *Trans. Amer. Math. Soc.* 274 (1982) 399–443.
- [K] O. Kerner, Wild hereditary algebras, in: Proc. Workshop at UNAM, Mexico, in: CMS Conf. Proc., vol. 19, 1996, pp. 65–108.
- [R] C.M. Ringel, *Tame Algebras and Integral Quadratic Forms*, Lecture Notes in Math., vol. 1099, Springer-Verlag, Heidelberg, 1984.
- [U] L. Unger, On the simplicial complex of tilting modules over quiver algebras, *Proc. London Math. Soc.* (3) 73 (1996) 27–46.